

The determinants

Mongi BLEL

King Saud University

July 4, 2019

Table of contents

- 1 Definition of The Determinant
- 2 Properties of the determinants
- 3 The adjoint matrix

Definition of The Determinant

Definition

If $A = (a_{j,k})$ is a square matrix of type n . Denote $A_{j,k}$ the square matrix of type $n - 1$ obtained by deleting the j^{th} -row and the k^{th} -column of A .

Example: If $A = \begin{pmatrix} 1 & 2 & 3 \\ -1 & 2 & -4 \\ 2 & -3 & 4 \end{pmatrix}$, then $A_{2,3} = \begin{pmatrix} 1 & 2 \\ 2 & -3 \end{pmatrix}$.

Definition

- ① If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, the determinant of A is defined by:

$$|A| = \det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$

- ② If $A = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$, the determinant of A is defined by:

$$|A| = \det(A) = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}.$$

Definition

3 If

$$A = \begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ a_{2,1} & a_{2,2} & \dots & a_{2,n} \\ & & \vdots & \\ a_{n,1} & a_{n,2} & \dots & a_{n,n} \end{pmatrix},$$

the determinant of A is defined by:

$$\begin{aligned} |A| = \det(A) &= a_{1,1}\det A_{1,1} + \dots + (-1)^{n+1}a_{1,n}\det A_{1,n} \\ &= \sum_{j=1}^n (-1)^{j+1}a_{1,j}\det A_{1,j}. \end{aligned}$$

Example

- ① If $A = \begin{pmatrix} 4 & 5 \\ 2 & 3 \end{pmatrix}$, the determinant of A is

$$|A| = \det(A) = \begin{vmatrix} 4 & 5 \\ 2 & 3 \end{vmatrix} = 4 \cdot 3 - 5 \cdot 2 = 2.$$

- ② If $A = \begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \\ 0 & 2 & 4 \end{pmatrix}$, the determinant of the matrix A is

$$|A| = \det(A) = \begin{vmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \\ 0 & 2 & 4 \end{vmatrix} = 1 \begin{vmatrix} 4 & 6 \\ 2 & 4 \end{vmatrix} - 3 \begin{vmatrix} 2 & 6 \\ 0 & 4 \end{vmatrix} + 5 \begin{vmatrix} 2 & 4 \\ 0 & 2 \end{vmatrix}$$

Example

3 If $A = \begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \\ 0 & 2 & 4 \end{pmatrix}$, the determinant of the matrix A is

$$|A| = \det(A) = \begin{vmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \\ 0 & 2 & 4 \end{vmatrix} = 1 \begin{vmatrix} 4 & 6 \\ 2 & 4 \end{vmatrix} - 3 \begin{vmatrix} 2 & 6 \\ 0 & 4 \end{vmatrix} + 5 \begin{vmatrix} 2 & 4 \\ 0 & 2 \end{vmatrix} = 0.$$

Definition

If A is a square matrix of order n , the determinant $\det A_{j,k}$ is called the minor of the entry $a_{j,k}$ or the $(j, k)^{\text{th}}$ minor of A and the number $C_{j,k} = (-1)^{j+k} \det A_{j,k}$ is called the cofactor of the entry $a_{j,k}$ or the $(j, k)^{\text{th}}$ cofactor of the matrix A .

Remark

- ① If A is a square matrix of order n , the determinant of the matrix A is equal to

$$\det A = \sum_{j=1}^n a_{1,j} C_{1,j}.$$

- ② By rearranging the boundaries we conclude to

$$\begin{aligned} \det A &= \sum_{j=1}^n a_{1,j} C_{1,j} = \sum_{j=1}^n a_{k,j} C_{k,j} \\ &= \sum_{k=1}^n a_{k,j} C_{k,j}. \end{aligned}$$

The Sarrus's Theorem

If $n = 3$ and the matrix

$$A = \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \\ a_{3,1} & a_{3,2} & a_{3,3} \end{pmatrix}$$

$$\begin{vmatrix} a_{1,1} & a_{1,2} & a_{1,3} & a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} & a_{2,3} & a_{2,1} & a_{2,2} \\ a_{3,1} & a_{3,2} & a_{3,3} & a_{3,1} & a_{3,2} \end{vmatrix}$$

$$\begin{aligned} \det A &= a_{1,1}(a_{2,2} \cdot a_{3,3} - a_{2,3} \cdot a_{3,2}) \\ &\quad - a_{1,2}(a_{2,1} \cdot a_{3,3} - a_{2,3} \cdot a_{3,1}) \\ &\quad + a_{1,3}(a_{2,1} \cdot a_{3,2} - a_{2,2} \cdot a_{3,1}) \end{aligned}$$

Example

$$\text{If } A = \begin{pmatrix} 3 & -4 & 0 \\ 0 & 7 & 6 \\ 2 & -6 & 1 \end{pmatrix},$$

$$\begin{aligned} \det A &= \begin{vmatrix} 3 & -4 & 0 & 3 & -4 \\ 0 & 7 & 6 & 0 & 7 \\ 2 & -6 & 1 & 2 & -6 \end{vmatrix} \\ &= 3 \cdot 7 \cdot 1 + (-4) \cdot 6 \cdot 2 - (-6) \cdot 6 \cdot 3 = 81. \end{aligned}$$

Properties of the Determinants

Theorem

- 1 If A is a square matrix, $\det A^T = \det A$.
- 2 If a square matrix A contains a zero row or column, then its determinant is 0.
- 3 If the matrix $A = (a_{j,k})_{1 \leq j,k \leq n}$ is upper (lower) triangular, then its determinant is equal to:

$$a_{1,1} \dots a_{n,n}.$$

- 4 If a square matrix A contains a row which is a multiple of a different row, then its determinant is 0.

Theorem

- 5 If a matrix B is obtained by multiplying a row of a matrix A by a number c , then $|B| = c|A|$ (i.e. $|c.R_j A| = c|A|$).
- 6 If a matrix B is obtained by interchanging two rows of a matrix A , then $\det B = -\det A$ (i.e. $|R_{j,k} A| = -|A|$).
- 7 If a matrix B is obtained by adding a multiple of a row to another row of a matrix A , then $\det B = \det A$. (i.e. $|cR_{j,k} A| = |A|$).

Example

$$\begin{aligned}
 & \begin{vmatrix} 1 & 3 & 2 & 2 \\ 2 & 3 & 3 & 1 \\ 3 & 3 & 4 & 2 \\ 1 & 1 & 1 & 1 \end{vmatrix} \begin{array}{l} (-2)R_{1,2}, (-3)R_{1,3} \\ \underline{\underline{\phantom{(-2)R_{1,2}, (-3)R_{1,3}}}} \\ (-1)R_{1,4} \end{array} \begin{vmatrix} 1 & 3 & 2 & 1 \\ 0 & -3 & -1 & -1 \\ 0 & -6 & -2 & -1 \\ 0 & -2 & -1 & 0 \end{vmatrix} \\
 & = \begin{array}{l} \phantom{(-1)R_{1,2}} \\ \phantom{(-1)R_{1,2}} \\ \phantom{(-1)R_{1,2}} \\ (-1)R_{1,2} \\ \underline{\underline{\phantom{(-1)R_{1,2}}}} \end{array} \begin{vmatrix} 3 & 1 & 1 \\ 6 & 2 & 1 \\ 2 & 1 & 0 \\ 3 & 1 & 1 \\ 3 & 1 & 0 \\ 2 & 1 & 0 \end{vmatrix} \\
 & = \begin{vmatrix} 3 & 1 \\ 2 & 1 \end{vmatrix} = -1.
 \end{aligned}$$

Example

$$\begin{vmatrix} 1 & -2 & 5 & -2 & -1 \\ -2 & 3 & -1 & 1 & 0 \\ 3 & -3 & 2 & 0 & -1 \\ 1 & -1 & 2 & 1 & -4 \\ 1 & -2 & 4 & -3 & 1 \end{vmatrix} \begin{matrix} 2R_{1,2}, -3R_{1,3} \\ = \\ -1R_{1,4}, -1R_{1,5} \end{matrix} \begin{vmatrix} 1 & -2 & 5 & -2 & -1 \\ 0 & -1 & 9 & -3 & -2 \\ 0 & 3 & -13 & 6 & 2 \\ 0 & 1 & -3 & 3 & -3 \\ 0 & 0 & -1 & -1 & 2 \end{vmatrix} \\
 = \begin{vmatrix} -1 & 9 & -3 & -2 \\ 3 & -13 & 6 & 2 \\ 1 & -3 & 3 & -3 \\ 0 & -1 & -1 & 2 \end{vmatrix} \\
 \begin{matrix} 3R_{1,2}, 1R_{1,3} \\ = \end{matrix} \begin{vmatrix} -1 & 9 & -3 & -2 \\ 0 & 14 & -3 & -4 \\ 0 & 6 & 0 & -5 \\ 0 & -1 & -1 & 2 \end{vmatrix}$$

$$\begin{aligned} &= \begin{vmatrix} 14 & -3 & -4 \\ 6 & 0 & -5 \\ 1 & 1 & -2 \end{vmatrix} \\ &= \begin{vmatrix} 14 & 6 & 1 \\ -3 & 0 & 1 \\ -4 & -5 & -2 \end{vmatrix} \\ (-1)\underline{\underline{R_{1,2}}}, 2\underline{\underline{R_{1,3}}} & \begin{vmatrix} 14 & 6 & 1 \\ -17 & -6 & 0 \\ 24 & 7 & 0 \end{vmatrix} \\ &= \begin{vmatrix} -17 & -6 \\ 24 & 7 \end{vmatrix} \\ \underline{\underline{1R_{1,2}}} & \begin{vmatrix} -17 & -6 \\ 7 & 1 \end{vmatrix} \\ &= 42 - 17 = 25. \end{aligned}$$

Example

$$\begin{vmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 2 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 2 & 1 & -2 \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 & 1 \\ 0 & -2 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & -3 \end{vmatrix} \\
 = \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & -3 \end{vmatrix} \\
 = \begin{vmatrix} 1 & 1 \\ 0 & -3 \end{vmatrix} = -3.$$

Example

$$\begin{aligned} \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} &= \begin{vmatrix} 1 & a & a^2 \\ 0 & b-a & (b-a)(b+a) \\ 0 & c-a & (c-a)(c+a) \end{vmatrix} \\ &= (b-a)(c-a) \begin{vmatrix} 1 & b+a \\ 1 & c+a \end{vmatrix} \\ &= (b-a)(c-a)(c-b). \end{aligned}$$

Theorem

If A and B are in $M_n(\mathbb{R})$, then

$$\det(AB) = \det A \det B.$$

Theorem

A square matrix A is invertible if and only if $\det A \neq 0$.

Remarks

- 1 If A is a square matrix of order n , then $|cA| = c^n|A|$.
- 2 Let A be a square matrix and B a row echelon form of A . Then there is a finite elementary matrices E_1, \dots, E_m such that $E_1 \dots E_m A = B$.

Moreover

$$\det(E_1) \dots \det(E_m) \det(A) = \det(B).$$

The adjoint matrix

Definition

Let A be a square matrix. The adjoint matrix associated to the matrix A is $\text{adj}(A) = (C_{j,k})^T$, where $(C_{j,k})$ is the cofactor matrix of A .

Theorem

Let A be a square matrix of order n , then

$$(\text{adj}(A))A = A(\text{adj}(A)) = (\det A)I_n.$$

Theorem

If A is an invertible matrix, then

$$A^{-1} = \frac{1}{\det A} \text{adj}(A).$$

$$\det A^{-1} = \frac{1}{\det A}.$$

Example

① $n = 2$, $A = \begin{pmatrix} 2 & -1 \\ 3 & 1 \end{pmatrix}$, $\det A = 5$, $\text{adj}(A) = \begin{pmatrix} 1 & 1 \\ -3 & 2 \end{pmatrix}$ and

$$A^{-1} = \frac{1}{5} \begin{pmatrix} 1 & 1 \\ -3 & 2 \end{pmatrix}.$$

② $n = 3$, $A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & -2 & 3 \\ -1 & 0 & 2 \end{pmatrix}$, $\det A = -13$

$$\text{adj}(A) = \begin{pmatrix} -4 & -5 & -2 \\ -2 & 4 & -1 \\ 3 & -6 & -5 \end{pmatrix}^T = \begin{pmatrix} -4 & -2 & 3 \\ -5 & 4 & -6 \\ -2 & -1 & -5 \end{pmatrix}$$

and $A^{-1} = \frac{1}{13} \begin{pmatrix} 4 & 2 & -3 \\ 5 & -4 & 6 \\ 2 & 1 & 5 \end{pmatrix}.$

Example

$$\textcircled{3} \quad n = 4, \quad A = \begin{pmatrix} 3 & 1 & 1 & 1 \\ 1 & 3 & 1 & 1 \\ 1 & 1 & 3 & 1 \\ 1 & 1 & 1 & 3 \end{pmatrix}, \quad \det A = -24$$

$$\text{adj}(A) = \begin{pmatrix} -20 & -4 & -4 & -4 \\ -4 & -20 & -4 & -4 \\ -4 & -4 & -20 & -4 \\ -4 & -4 & -4 & -20 \end{pmatrix},$$

$$\text{and } A^{-1} = \frac{1}{6} \begin{pmatrix} 5 & 1 & 1 & 1 \\ 1 & 5 & 1 & 1 \\ 1 & 1 & 5 & 1 \\ 1 & 1 & 1 & 5 \end{pmatrix}$$

Exercises

Let A be the following matrix $A = \begin{pmatrix} 1 & -4 & 3 \\ 0 & 2 & -1 \\ 1 & -2 & 3 \end{pmatrix}$.

- 1 Find the matrix $\text{adj}(A)$ and the determinant of the matrix A .
- 2 Find the inverse of the matrix A if it exists.

Example

Let A, B be matrices of size $(3, 3)$ such that A is not invertible and $|B| = 2$.

Find $|A \operatorname{adj}(A) + 2B^{-1}|$.

$$|A \operatorname{adj}(A) + 2B^{-1}| = \frac{8}{|B|} = 4.$$

Exercises

Let A and B be the following matrices:

$$A = \begin{pmatrix} 2 & 0 & 1 \\ 0 & 2 & 1 \\ 2 & -1 & 0 \end{pmatrix} \quad B = \begin{pmatrix} 3 & -1 & -1 \\ 2 & 0 & -1 \\ -2 & 1 & 4 \end{pmatrix}$$

Find the number a such that $A^2 - AB + aI_3 = 0$ and find the inverse matrix of A .

If $\text{adj}(A) = \begin{pmatrix} 3 & 1 & 2 \\ 0 & 3 & 2 \\ -1 & 2 & 1 \end{pmatrix}$, find the matrix A .

We have $A\text{adj}(A) = I_3$, then $A = |A|(\text{adj}(A))^{-1}$ and $|A| |\text{adj}(A)| = |A|^3$, then $|\text{adj}(A)| = |A|^2$. Therefore $A = \sqrt{|\text{adj}(A)|}(\text{adj}(A))^{-1}$.

$|\text{adj}(A)| = 1$, the $|A| = 1$ and $A = \begin{pmatrix} -1 & 3 & -4 \\ -2 & 5 & -6 \\ 3 & -7 & 9 \end{pmatrix}$

Exercises

- ① Prove that if a matrix $A \in M_n(\mathbb{R})$, $n \geq 2$ has an inverse, then

$$\text{adj}(\text{adj}(A)) = (\det A)^{n-2}A.$$

- ② Prove that a matrix A has an inverse if and only if the matrix $\text{adj}(A)$ has an inverse.

Solution

- ① From the relation $A \operatorname{adj}(A) = |A|I_n$ we conclude that $|\operatorname{adj}(A)| = |A|^{n-1}$ and $(\operatorname{adj}(A))^{-1} = \frac{1}{|A|}A$ if $|A| \neq 0$.

If the matrix A has an inverse, then $A^{-1} = \frac{1}{|A|} \operatorname{adj}(A)$.

Let $B = \operatorname{adj}(A)$, then $B \operatorname{adj}(B) = |B|I_n = |A|^{n-1}I_n$ and $\operatorname{adj}(B) = |A|^{n-1}B^{-1} = |A|^{n-2}A$.
 $\operatorname{adj}(\operatorname{adj}(A)) = (\det A)^{n-2}A$.

- ② From the relation $A \operatorname{adj}(A) = |A|I_n$ we conclude that if the matrix A has an inverse then the matrix $\operatorname{adj}(A)$ has an inverse. Also from the same relation, if the matrix $\operatorname{adj}(A)$ has an inverse and the matrix A do not has an inverse, then $A \operatorname{adj}(A) = 0$ and $A \operatorname{adj}(A)(\operatorname{adj}(A))^{-1} = 0$. Then $A = 0$ this is absurd, because if $A = 0$ then $\operatorname{adj}(A) = 0$.